

A NOTE ON MULTIPLE DIRICHLET'S q - L -FUNCTION

TAEKYUN KIM

*Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr*

ABSTRACT. Recently, the two variable q - L -functions which interpolate the generalized q -Bernoulli polynomials associated with χ are introduced and studied, cf. [2]. In this paper, we construct multiple Dirichlet's q - L -function which interpolates the generalized multiple q -Bernoulli numbers attached to χ at negative integer and study its properties.

§1. INTRODUCTION

Let $0 < q < 1$ be and for any positive integer k , define its q -analogue $[k]_q = \frac{1-q^k}{1-q}$. Let $\mathbb{C}$ be the field of complex numbers. The q -Bernoulli numbers are usually defined as

$$(1) \quad \beta_{0,q} = \frac{q-1}{\log q}, \quad (q\beta_q + 1)^n - \beta_{n,q} = \delta_{n,1},$$

where $\delta_{n,1}$ is Kronecker symbol and we use the usual convention about replacing β_q^i by $\beta_{i,q}$, cf. [2, 4, 5, 6]. Note that $\lim_{q \rightarrow 1} \beta_{k,q} = B_k$, where B_k are the k -th ordinary Bernoulli numbers. In [3, 5, 6], the q -Bernoulli polynomials are also defined by

$$(2) \quad F_q(t, x) = \frac{q-1}{\log q} e^{\frac{t}{1-q}} - t \sum_{n=0}^{\infty} q^{n+x} e^{[n+x]_q t} = \sum_{n=0}^{\infty} \frac{\beta_{n,q}(x)}{n!} t^n, \text{ for } x \in \mathbb{C}.$$

Key words and phrases. zeta function, L -function, Bernoulli numbers.
2000 Mathematics Subject Classification: 11S80, 11B68, 11M99 .

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

From (1) and (2), we can derive the below formula:

$$\beta_{n,q}(x) = \sum_{i=0}^n \binom{n}{i} q^{xi} \beta_{i,q}[x]_q^{n-i} = \left(\frac{1}{1-q}\right)^n \sum_{i=0}^n \binom{n}{i} (-1)^i q^{xi} \frac{i}{[i]_q},$$

where $\binom{n}{i}$ is the binomial coefficient. In the recent paper the multiple generalized Bernoulli numbers attached to χ , $B_{n,\chi}^{(r)}$, are defined by

(3)

$$F_{\chi}^r(t) = \sum_{a_1, \dots, a_r=1}^f \frac{\chi(\sum_{i=1}^r a_i) t^r e^{(\sum_{i=1}^r a_i)t}}{(e^{ft} - 1)^r} = \sum_{n=0}^{\infty} B_{n,\chi}^{(r)} \frac{t^n}{n!}, \text{ for } |t| < 2\pi/f, \text{ (see [1])}.$$

In this paper we consider the q -analogue of the above multiple generalized Bernoulli numbers attached to χ . Finally we construct the multiple q - L -function which interpolates the q -analogue of multiple generalized Bernoulli numbers attached to χ at negative integers and investigate its properties.

2. MULTIPLE DIRICHLET'S q - L -FUNCTION

Let $\chi(\neq 1)$ be the primitive Dirichlet's character with conductor $f \in \mathbb{N}$. Then the generalized q -Bernoulli numbers attached to χ are defined as

$$(4) \quad F_{q,\chi}(t) = -t \sum_{a=1}^f \chi(a) \sum_{n=0}^{\infty} q^{fn+a} e^{[fn+a]_q t} = \sum_{n=0}^{\infty} \beta_{n,\chi,q} \frac{t^n}{n!}, \text{ cf. [5].}$$

By (2) and (4), we easily see that

$$\beta_{n,\chi,q} = [f]_q^{n-1} \sum_{a=1}^f \chi(a) \beta_{n,q^f}\left(\frac{a}{f}\right).$$

By the meaning of Eq.(4), we can consider the multiple generalized q -Bernoulli numbers attached to χ as follows:

$$(5) \quad \begin{aligned} F_{q,\chi}^r(t) &= (-t)^r \sum_{a_1, \dots, a_r=1}^f \chi\left(\sum_{i=1}^r a_i\right) \sum_{n_1, \dots, n_r=0}^{\infty} q^{\sum_{i=1}^r (a_i + n_i f)} e^{([\sum_{i=1}^r (a_i + n_i f)]_q t} \\ &= \sum_{n=0}^{\infty} \beta_{n,\chi,q}^{(r)} \frac{t^n}{n!}. \end{aligned}$$

Note that $\lim_{q \rightarrow 1} \beta_{n,\chi,q}^{(r)} = B_{n,\chi}^{(r)}$. Let $\Gamma(s)$ be the gamma function. Then it is easy to see that

$$\begin{aligned}
 & \frac{1}{\Gamma(s)} \int_0^\infty F_{\chi,q}^r(-t) t^{s-1-r} dt \\
 (6) \quad &= \sum_{n_1, \dots, n_r=1}^\infty \chi(n_1 + \dots + n_r) q^{n_1 + \dots + n_r} \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-[n_1 + \dots + n_r]_q t} dt \\
 &= \sum_{n_1, \dots, n_r=1}^\infty \frac{\chi(n_1 + \dots + n_r)}{[n_1 + \dots + n_r]_q^s} q^{n_1 + \dots + n_r}.
 \end{aligned}$$

Thus, we can define the multiple Dirichlet's q - L -function as follows:

$$(7) \quad L_q^r(s, \chi) = \sum_{n_1, \dots, n_r=1}^\infty q^{n_1 + \dots + n_r} \frac{\chi(n_1 + \dots + n_r)}{[n_1 + \dots + n_r]_q^s}, \text{ for } s \in \mathbb{C},$$

where χ is non-trivial primitive Dirichlet's character with conductor $f \in \mathbb{N}$. By (5), (6) and (7), we easily see that

$$(8) \quad L_q^r(-n, \chi) = (-1)^r \frac{n!}{(n+r)!} \beta_{n+r,\chi,q}^{(r)}, \text{ for } n \in \mathbb{N}.$$

Let s be a complex variable, a and F be integers with $0 < a < F$. We now consider the function $H_{r,q}(s; a_1, \dots, a_r | F)$ as follows:

$$(9) \quad H_{r,q}(s; a_1, \dots, a_r | F) = \sum_{\substack{m_1, \dots, m_r > 0 \\ m_i \equiv a_i \pmod{F}}} \frac{q^{m_1 + \dots + m_r}}{[m_1 + \dots + m_r]_q^s} = [F]_q^{-s} \zeta_{r,q^F}(s, \frac{a_1 + \dots + a_r}{F}),$$

where $\zeta_{r,q}(s, a) = \sum_{n_1, \dots, n_r=0}^\infty \frac{q^{n_1 + \dots + n_r + a}}{[n_1 + \dots + n_r + a]_q^s}$, (see [5, 6]).

The function $H_{r,q}(s; a_1, \dots, a_r | F)$ is a meromorphic for $s \in \mathbb{C}$ with poles at $s = 1, \dots, r$. In [5], the multiple q -Bernoulli polynomials are defined by

$$(10) \quad (-t)^r \sum_{n_1, \dots, n_r=0}^\infty q^{x + \sum_{i=1}^r n_i} e^{[x + \sum_{i=1}^r n_i]_q t} = \sum_{n=0}^\infty B_{n,q}^{(r)}(x) \frac{t^n}{n!}.$$

For $n \in \mathbb{N}$, it was well known that

$$(11) \quad \zeta_{r,q}(-n, x) = (-1)^r \frac{n!}{(n+r)!} B_{n+r,q}^{(r)}(x), \text{ cf. [5]}.$$

Let $F \in \mathbb{N}$ be the conductor of χ . Then the multiple Dirichlet's q - L -function can be expressed as the sum

$$(12) \quad L_q^r(s, \chi) = \sum_{a_1, \dots, a_r=1}^F \chi(a_1 + \dots + a_r) H_{r,q}(s; a_1, \dots, a_r | F), \text{ for } s \in \mathbb{C}.$$

By (9) and (12), we easily see that

$$(13) \quad H_{r,q}(-n; a_1, \dots, a_r | F) = [F]_q^n (-1)^r \frac{n!}{(n+r)!} B_{n+r, q^F}^{(r)}\left(\frac{a_1 + \dots + a_r}{F}\right), \text{ for } n \in \mathbb{N}.$$

From (5) and (10), we can also derive the below formula:

$$(14) \quad \beta_{n, \chi, q}^{(r)} = [F]_q^n \sum_{a_1, \dots, a_r=1}^F \chi(a_1 + \dots + a_r) B_{n, q^F}^{(r)}\left(\frac{a_1 + \dots + a_r}{F}\right).$$

By using (12), (13) and (14), we easily see that

$$L_q^r(-n, \chi) = (-1)^r \frac{n!}{(n+r)!} \beta_{n+r, \chi, q}^{(r)}.$$

In Eq.(10), we note that

$$(15) \quad B_{n,q}^{(r)}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{xk} B_{k,q}^{(r)}, \text{ cf. [5, 6]},$$

where $B_{n,q}^{(r)} = B_{n,q}^{(r)}(0)$. From Eq.(15), we derive the function $H_{r,q}(s; a_1, \dots, a_r | F)$ which is modified by

$$(16) \quad \begin{aligned} & H_{r,q}(s; a_1, \dots, a_r | F) \\ &= \frac{1}{[F]_q^r} \frac{[\sum_{i=1}^r a_i]_q^{-s+r}}{\prod_{j=1}^r (s-j)} \sum_{k=0}^{\infty} \binom{-s+r}{k} \left(\frac{[F]_q}{[\sum_{i=1}^r a_i]_q} \right)^k q^{(\sum_{i=1}^r a_i)k} B_{k,q}^{(r)}. \end{aligned}$$

By (12) and (16), we obtain the following:

$$(17) \quad \begin{aligned} L_q^r(s, \chi) &= \frac{1}{\prod_{j=1}^r (s-j)} \frac{1}{[F]_q^r} \sum_{a_1, \dots, a_r=1}^F \chi(a_1 + \dots + a_r) [a_1 + \dots + a_r]_q^{-s+r} \\ &\quad \cdot \sum_{m=0}^{\infty} \binom{r-s}{m} \left(\frac{[F]_q}{[a_1 + \dots + a_r]_q} \right)^m q^{(a_1 + \dots + a_r)m} B_{m,q}^{(r)}. \end{aligned}$$

Finally, we suggest the below question

Question. *Is it possible to give the p -adic analogue of Eq.(17) which can be viewed as interpolating, in the same way that $L_{p,q}(s, \chi)$ interpolates $L_q(s, \chi)$ in [2, 7] ?*

REFERENCES

1. L.C. Jang, T. Kim, D.W. Park, *Kummer congruences for the Bernoulli numbers of higher order*, Appl. Math. Comput. **151** (2004), 589-593.
2. T. Kim, *Power series and asymptotic series associated with the q -analogue of two variable p -adic L -function*, Russ. J. Math. Phys. **12** (2005), 186-196.
3. T. Kim, L.C. Jang, S.H. Rim, *An extension of q -zeta function*, Int. J. Math. Math. Sci. **2004 no. 49** (2004), 2649-2651.
4. T. Kim, *On multiple L -function*, Proceedings of the Japan-Korea Joint Seminar on Number Theory (edited by Taguchi and Kim) **1 no. 1** (2004), 110-116.
5. T. Kim, *Analytic continuation of multiple q -zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11** (2004), 71-76.
6. T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9 no.3** (2002), 288-299.
7. T. Kim, *Multiple p -adic L -functions*, Russ. J. Math. Phys. **13 no.1** (2006), 00-00.